



Polymer Models of Yeast *S. cerevisiae* Genome Organization

Citation

Fudenberg, Geoff, Jon-Matthew Belton, Anton Goloborodko, Maxim Imakaev, Job Dekker, and Leonid Mirny. 2013. Polymer models of yeast *s. cerevisiae* genome organization. *Epigenetics & Chromatin* 6(Suppl 1): P128.

Published Version

doi:10.1186/1756-8935-6-S1-P128

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:11732107>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)



POSTER PRESENTATION

Open Access

Polymer models of yeast *S. cerevisiae* genome organization

Geoff Fudenberg^{1*}, Jon-Matthew Belton², Anton Goloborodko³, Maxim Imakaev³, Job Dekker², Leonid Mirny^{3,4}

From Epigenetics & Chromatin: Interactions and processes
Boston, MA, USA. 11-13 March 2013

Introduction

Three-dimensional (3D) chromosomal organization impacts critical cellular processes including transcription, replication, and genomic stability. Despite the ubiquity of these challenges, recent 3C-based experiments [1] suggest that major features of interphase chromosomal organization vary across eukaryotes. In human cells, Hi-C revealed that chromosomal regions of similar functional state (eg. high gene expression) were enriched for contact probability in 3D [2]. Conversely, 3D contact probability was depleted between regions with different functional states. Moreover, the probability of a contact between two genomic loci was found to be inversely proportional to intervening genomic distance, s^{-1} and consistent with an unknotted non-equilibrium fractal globular polymer state. Here we analyze new, high-coverage, Hi-C data and developed stochastic simulations of polymer dynamics to study the spatial organization of yeast chromosomes.

Results

Unlike in human chromosomes, we find no evidence of a domain-type organization in yeast. Furthermore, we find that contact probability decays more like $s^{-3/2}$ with increasing genomic distance in *cerevisiae*; this indicates that yeast chromosomes do not exhibit a fractal globule organization. Instead, we find that a Rabl-like organization of chromosomes and constraints from the nucleolus appear to be the most prominent features of chromatin organization [3,4]. In this Rabl organization, centromeres are co-localized near the spindle pole body on one side of the nucleus. Our stochastic polymer simulations allow us to reconstruct conformational ensembles consistent with Hi-C maps. We demonstrate that a Rabl organization leads to cross-like patterns of interactions between centromeric regions, as observed in yeast Hi-C data. Our simulations also allow us

to match experimental data on diffusion of individual genomic loci; this allows us to study the temporal evolution of chromosomal conformations.

Conclusions

Our models show that: (1) yeast chromosomes are generally consistent with a mildly confined “equilibrium globular” polymer state, in contrast with observations in human cell lines; (2) tethering in a Rabl conformation induces a ‘polymer brush’ effect which reproduces the majority of observed intra- and inter- chromosomal Hi-C interactions in yeast; (3) rapid progression through the cell cycle allows for spatial, but not necessarily topological, equilibration of yeast chromosomes, limiting their mutual entanglement.

Author details

¹Biophysics, Harvard University, Boston, MA, 02115, USA. ²Systems Biology, UMMS Medical School, Worcester, MA, 01605, USA. ³Physics, MIT, Cambridge, MA, 02139, USA. ⁴Division of Health Science and Technology, Harvard-MIT, Cambridge, MA, 02139, USA.

Published: 8 April 2013

References

1. Dekker J, Rippe K, Dekker M, Kleckner N: **Capturing Chromosome Conformation.** *Science* 2002, **295**:1306.
2. Lieberman-Aiden E, van Berkum NL, Williams L, Imakaev M, Ragoczy T, Telling A, Amit I, Lajoie BR, Sabo PJ, Dorschner MO, Sandstrom R, Bernstein B, Bender MA, Groudine M, Gnirke A, Stamatoyannopoulos J, Mirny LA, Lander ES, Dekker J: **Comprehensive Mapping of Long-Range Interactions Reveals Folding Principles of the Human Genome.** *Science* 2009, **326**:289.
3. Duan Z, Andronescu M, Schutz K, Mcllwain S, Kim YJ, Lee C, Shendure J, Fields S, Blau CA, Noble WS: **A three-dimensional model of the yeast genome.** *Nature* 2010, **465**(7296):363-367.
4. Therizols P, Duong T, Dujon B, Zimmer C, Fabre E: **Chromosome arm length and nuclear constraints determine the dynamic relationship of yeast subtelomeres.** *Proc Natl Acad Sci* 2010, **107**(5):2025-2030.

doi:10.1186/1756-8935-6-S1-P128

Cite this article as: Fudenberg et al.: Polymer models of yeast *S. cerevisiae* genome organization. *Epigenetics & Chromatin* 2013 **6**(Suppl 1):P128.

¹Biophysics, Harvard University, Boston, MA, 02115, USA
Full list of author information is available at the end of the article